

#6472 Store at -20°C

# SignalSilence® FAK siRNA I



✓ 10 µM in 300 µl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com  
Support ■ 877-678-TECH (8324) info@cellsignal.com  
Web ■ www.cellsignal.com

rev. 02/11/16

For Research Use Only. Not For Use In Diagnostic Procedures.


### Species Cross-Reactivity: H

**Description:** SignalSilence® FAK siRNA I from Cell Signaling Technology (CST) allows the researcher to specifically inhibit FAK expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

**Background:** Focal adhesion kinase (FAK) is a widely expressed cytoplasmic protein tyrosine kinase involved in integrin-mediated signal transduction. It plays an important role in the control of several biological processes, including cell spreading, migration, and survival (1). Activation of FAK by integrin clustering leads to autophosphorylation at Tyr397, which is a binding site for the Src family kinases PI3K and PLCγ (2-5). Recruitment of Src family kinases results in the phosphorylation of tyrosine residues 407, 576, and 577 in the catalytic domain, and tyrosine residues 871 and 925 in the carboxy-terminal region of FAK (6,7).

**Directions for Use:** CST recommends transfection with 100 nM FAK siRNA I 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

**Quality Control:** Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from HeLa cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® FAK siRNA I (+) or SignalSilence® FAK siRNA II #6483 (+), using FAK Antibody #3285 (upper) or α-Tubulin (11H10) Rabbit mAb #2125 (lower). The FAK Antibody confirms silencing of FAK expression, while the α-Tubulin (11H10) Rabbit mAb is used as a loading control.

Entrez-Gene ID #5747  
Swiss-Prot Acc. #Q05397

**Storage:** FAK siRNA I is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit [www.cellsignal.com](http://www.cellsignal.com) for a complete listing of recommended companion products.

### Background References:

- (1) Parsons, J.T. et al. (2000) *Oncogene* 19, 5606-5613.
- (2) Schaller, M.D. et al. (1994) *Mol. Cell. Biol.* 14, 1680-1688.
- (3) Cobb, B.S. et al. (1994) *Mol. Cell. Biol.* 14, 147-155.
- (4) Chen, H.C. et al. (1996) *J. Biol. Chem.* 271, 26329-26334.
- (5) Zhang, X. et al. (1999) *Proc. Natl. Acad. Sci. USA* 96, 9021-9026.
- (6) Calalb, M.B. et al. (1995) *Mol. Cell. Biol.* 15, 954-963.
- (7) Schlaepfer, D.D. et al. (1994) *Nature* 372, 786-791.

© 2011 Cell Signaling Technology, Inc. SignalSilence® CST™ and Cell Signaling Technology® are trademarks of Cell Signaling Technology, Inc.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.